Can we observe the σ meson in peripheral heavy ion collisions?

A. A. Natale and C. G. Roldão
Instituto de Física Teórica, Universidade Estadual Paulista, Rua Pamplona 145, 01405-900, São Paulo, SP, Brazil

(February 8, 2008)

We discuss the possible production of the σ meson and detection of its two photon decay mode in peripheral heavy ion collisions at RHIC. Assuming the mass and total decay width values reported recently by the E791 Collaboration, we verify that the σ meson can be observed in the subprocess $\gamma\gamma \to \gamma\gamma$ if its partial decay width into photons is of the order of a few keV.

arXiv:nucl-th/0204004v2 2 Apr 2002

The possible existence of a light scalar sigma (σ) meson has been a controversial subject for roughly forty years [1]. In the strong interaction phenomena the σ meson plays a role analogous to the Higgs particle in the Weinberg-Salam theory. It is expected that when the chiral symmetry of the strong force between quarks is broken by the nonperturbative vacuum, generating a massless pseudoscalar meson (the pion), a massive scalar meson (the σ) is also generated [2]. The sigma meson is fundamental for effective theories describing the nuclear force, where it is responsible for a attractive part of the nuclear potential [3]. Although its experimental confirmation would be quite important, its existence has remained inconclusive for a long time making it disappear from the review of particle physics and reappearing only recently [1,4]. There are two difficulties for its discovery: the extraction of the scalar properties from experiment and the lack of knowledge about its underlying quark substructure. Because the $J = 0$ channel may contain strong competing contributions, such resonance may not necessarily dominate its amplitudes and could be hard to "observe". In such an instance its verification would be linked to the model used to its description.

One of the most compelling evidences for the σ meson was reported recently by the E791 Collaboration, resulting from the observation of the $D^+ \to \pi^-\pi^+\pi^+$ decay [5]. The Dalitz-plot of this decay can hardly be fitted without a 0^{++} (σ) resonance, whose mass and total decay width are $478^{+24}_{-23} \pm 17$ MeV and $324^{+42}_{-40} \pm 21$ MeV, respectively. The σ meson appears as an intermediate state decaying into two pions, which is the same final state questioned over many years as a possible (or not) signal for this broad resonance.

The σ meson can also decay into two photons, although this decay is orders of magnitude smaller than the $\pi\pi$ decay mode it has the advantage that the final state is not strongly interacting. Therefore, the observation of the decay $\sigma \to \gamma\gamma$ would be a cleaner way to establish its existence. This decay is also important to unravel any possible amount of mixing of quarks or gluons in this resonance [6]. The ideal observation of this resonance would be through the process $\gamma\gamma \to \sigma \to \gamma\gamma$, where the production and decay modes are free of any problem caused by the strong interaction of the initial or final states which may hide its properties. Our work will focus exactly on the possibility of observing this meson in peripheral heavy ion collisions through the subprocess $\gamma\gamma \to \gamma\gamma$.

In the Relativistic Heavy-Ion Collider (RHIC), operating at the Brookhaven Laboratory, there is a program for very peripheral heavy ion collisions, where each charge Z accelerated ion has a strong photon field around it that may originate electromagnetic interactions, and where the ions remain intact after these interactions. The huge luminosity of photons in these collisions open the possibilities of studying two-photon physics as discussed by several authors [7], and, in particular, the process $\gamma\gamma \to \gamma\gamma$ could indeed be measured [8].

We compute the $\gamma\gamma \to \gamma\gamma$ subprocess, for the Au ion at RHIC energies, $\sqrt{s} = 200$ GeV/nucleon. The photon distribution in the nucleus can be described using the equivalent-photon or Weizsäcker-Williams approximation in the impact parameter space. We use the photon distribution obtained by Cahn and Jackson [9], wich takes into account the condition for realistic peripheral collisions (for details see Ref. [10]). The subprocess $\gamma\gamma \to \gamma\gamma$ up to energies of a few GeV is dominated by the continuous fermion box diagram, and is a background for the resonant $\gamma\gamma \to \sigma \to \gamma\gamma$ process. It was first calculated exactly by Karplus and Neuman [11] and De Tollis [12]. There are sixteen helicity amplitudes for the process and, due to symmetry properties, the number of independent amplitudes will be only five, that may be chosen to be M_{++++}, M_{++--}, M_{+-+-}, M_{+--+} and M_{+++-}. Where the + or − denotes the circular polarization values +1 and −1. The remaining helicity amplitudes may be obtained from parity and permutation symmetry. Of these five helicity amplitudes, three are related by crossing, hence it is sufficient to give just three, which are presented in detail in Ref. [10].

The electron gives the major contribution to the continuous $\gamma\gamma \to \gamma\gamma$ subprocess. The muon and u contributions are smaller roughly by one order of magnitude, and the d and s quark contributions are even smaller. The process is proportional to $(q_f^2)^4$ where q_f is their charge, and this is the main reason for the suppression of the quarks contribution [10]. Contribution from pion loops, heavy quarks or leptons, and charged weak bosons are negligible up to O(2 GeV). The main background for this reaction could come from the processes: a) Double-Pomeron exchange, this process was computed and after extracting the contribution of central collisions (see Ref. [13] for details of the procedure) they are shown to be negligible, in agreement with the conclusions of our work about the importance of double Pomeron contributions to ultra peripheral heavy ion collisions [13], b) Double bremsstrahlung (which dominates the region of $|\cos\theta| \approx 1$, where θ is the scattering angle in the two photon center-of-mass system). This background is eliminated with cuts in the scattering angle, and in all calculations we will consider the maximum value $|\cos\theta| = 0.5$. This cut is also consistent with all the requirements proposed by Nystrand and Klein [14] to eliminate background, and retains only the back-to-back photons in the central region of rapidity. However, a full simulation of bremsstrahlung contributions should be performed to estimate the effectiveness of this cut. Note that the angular cut diminishes the contribution of the continuous process (peaked at large angles) in comparison with the isotropic photon distribution produced by a scalar resonance. As discussed in Ref. [8], the $\gamma\gamma$ scattering can indeed be measured in peripheral heavy ion collisions. Just to give one idea of the number of events, with a luminosity of 2.0 $\times 10^{26}$ cm^{-2}s^{-1} [14] and integrating in a bin of energy 700 ± 100 MeV (which is free of any strong resonance decaying into two-photons), we have 1532 events/year assuming 100% efficiency in the tagging of the ions and photon detection [10].

Another possible background is the continuous or resonant (σ) production of a pair of neutral pions where both are misidentified with photons. This accidental background can be easily isolated measuring its invariant mass distribution, and making a cut that discriminates a single photon from one pion that subsequently decays into two photons. For example, in the case of the sigma meson each one of the neutral pions from its much large hadronic decay should be misidentified. These pions would produce pairs of photons with a large opening angle ϕ, where $\cos(\phi/2) = \sqrt{1 - 4m_\pi^2/m_\sigma^2}$. However, the calorimeters already in use in many experiments are able to distinguish between these two and single photon events with high efficiency (see, for instance, Ref. [16]). The detectors at RHIC are prepared to detect photons of O(1) GeV, because these are also a signal for the quark-gluon plasma formed in central collisions.

The contribution from $\gamma-$Pomeron$\rightarrow$ V $\rightarrow \gamma\gamma X$ may also produce another accidental background. The vector mesons, V, are produced with a p_T distribution similar to the σ and at higher rates than $\gamma\gamma \rightarrow$ scalar/tensor, *e.g.* $\gamma-$Pomeron$\rightarrow \omega$ rate is 10 Hz at RHIC [15], three orders of magnitude higher than for a similar mass meson of spin 0 or 2. The ω branching ratio to three photons is 8.5%. If a small p_T photon from this decay is undetected, one is left with a low p_T two-photon final state with $M < 780$ MeV that could be taken for a σ candidate. This background is likely to be much greater than the proposed signal. Furthermore, the σ is very wide, making background subtraction difficult. At higher masses, one also has $\phi \rightarrow \eta\gamma$, $\pi^0\gamma$, $K_L K_S \rightarrow \gamma X$ as well as $\gamma\gamma \rightarrow f_2(1270) \rightarrow \pi^0\pi^0$ and possibly copious production of $\rho(1450)$ and $\rho(1700)$ by $\gamma-$Pomeron interactions. Clearly a full simulation of all these background processes should be kept in mind when measuring two-photon final states.

Photon pair production via the box diagram is a background to $\gamma\gamma \rightarrow \sigma \rightarrow \gamma\gamma$ process (or vice versa), both have the same initial and final states, and for this reason they can interfere one in another. Normally the interference between a resonance (R) and a continuum process is unimportant, because on resonance the two are out of phase. Assuming a Breit-Wigner profile the total cross section for the elementary subprocess $\gamma\gamma \rightarrow R \rightarrow \gamma\gamma$ is

$$\frac{d\sigma_{ZZ}^{\gamma\gamma}}{dM} = 16\pi \frac{dL}{dM} \frac{\Gamma^2(R \rightarrow \gamma\gamma)}{(M^2 - m_R^2)^2 + m_R^2 \Gamma_{total}^2}, \qquad (1)$$

where L is the photon luminosity, M is the energy of the photons created by the collision of the ions. $\Gamma(R \rightarrow \gamma\gamma)(\equiv \Gamma_{\gamma\gamma})$ and Γ_{total} are the partial and total decay width of the resonance with mass m_R in its rest frame. The results for the box diagram and the σ production in the Breit-Wigner approximation are shown in Fig.(1). We used the value $\Gamma_{\gamma\gamma} = 4.7$ keV determined by Pennington and Boglione [17]. This value is in the middle of various determinations (for other values see Ref. [10]). The σ mass and total decay width correspond to the central values of Ref. [5]. The use of a constant total width in the σ resonance shape is not always a good approximation [18]. We discuss the $\gamma\gamma \rightarrow \gamma\gamma$ process above the two pions threshold where the peculiarities of the broad resonance, basically due to the σ decay into pions, are not so important. However, we also computed the cross section with a energy dependent total width $\Gamma(M) \simeq \Gamma_0 \left(p^*/p_0^*\right)^{2J+1}$, where $p^* = p^*(M)$ is the momentum of decay particles at mass M, measured in the resonance rest frame, $p_0^* = p^*(m_R)$, which, as shown by Jackson many years ago [19], is more appropriate for a quite broad resonance. The net effect is a slight distortion of the cross section shape with a small increase of the total cross section, and we shall not consider it again.

Off resonance we can expect a negligible contribution for the process $\gamma\gamma \rightarrow R \rightarrow \gamma\gamma$ and consequently the same for its interference with the continuum process. This is true if the resonance has a small total decay width [10], but this is not the σ case. To take into account the interference we must make use of a model to calculate the helicity amplitudes of the σ meson exchange. Using the effective lagrangian $g_s F^{\mu\nu} F_{\mu\nu} \Phi_s$, where g_s is the coupling of the photons to the scalar field Φ_s and $F^{\mu\nu}$ is the electromagnetic field tensor the following amplitudes comes out [10,20]:

$$M_{++++} = -\frac{2\pi}{\alpha^2} F(\lambda r_t),$$

$$
\begin{aligned}
M_{+-+-} &= -\frac{2\pi}{\alpha^2}F(\lambda t_t),\\
M_{+--+} &= -\frac{2\pi}{\alpha^2}F(\lambda s_t),\\
M_{+++-} &= 0,\\
M_{++--} &= -\frac{2\pi}{\alpha^2}\{F(\lambda s_t)+F(\lambda t_t)+F(\lambda r_t)\}. \qquad (2)
\end{aligned}
$$

where α is the fine-structure constant, $\lambda = (m_f/m_R)^2$, and

$$
F(x) = 16x^2\frac{\Gamma_{\gamma\gamma}}{m_R}\left(4x - 1 + i\frac{\Gamma_{total}}{m_R}\right)^{-1}, \qquad (3)
$$

and r_t, s_t and t_t are related with the standard Mandelstam variables s, t, and u by $r_t = \frac{1}{4}\frac{s}{m_f^2}$, $s_t = \frac{1}{4}\frac{t}{m_f^2}$ and $t_t = \frac{1}{4}\frac{u}{m_f^2}$.

These amplitudes and the ones describing the fermion (with mass m_f) box diagram enter in the expression for the differential cross section of photon pair production from photon fusion $d\sigma/d\cos\theta = (1/2\pi)(\alpha^4/s)(\sum_f q_f^2)^4 \sum |M|^2$, to give the total cross section of Fig.(1). We verified that the interference is destructive. The continuum process ($PP \to \gamma\gamma$) is also shown in Fig.(1). The diffractive resonant event is even smaller and is not shown in this figure.



FIG. 1. Invariant mass distribution of photon pair production. The solid curve is due to box diagram only, the dashed one is due to the process $\gamma\gamma \to \sigma \to \gamma\gamma$ in the Breit-Wigner approximation, the dash-dotted is the scalar contribution of Eq.(2), and the dotted one is the continuum Pomeron process $PP \to \gamma\gamma$. We assumed $\Gamma_{\gamma\gamma} = 4.7$ keV, and the angular cut is equal to $-0.5 < \cos\theta < 0.5$.

The effective Lagrangian model used to compute the σ contribution to the photon pair production gives a larger cross section than the calculation with the Breit-Wigner approximation at energies above $M \approx 600$ MeV. It is dominated by the s channel contribution. We consider the Breit-Wigner result as the best signal representation for the resonant process because we are using the E791 data and this one was fitted by a Breit-Wigner profile. The effective Lagrangian gives a nonunitary amplitude that overestimates the sigma production above 600 MeV and shows the model dependence in the σ analysis that we commented before. The Breit-Wigner profile is not a bad approximation as long as we stay above the two pions threshold and in the following we assume that the signal is giving by it (the dashed curve of Fig.(1)) and the background is giving by the box diagram result (the solid curve of Fig.(1)). Note that, due to the destructive interference, the actual measurement will give a curve below the solid curve of Fig.(1).

From the experimental point of view we would say that the reaction $\gamma\gamma \to \gamma\gamma$ has to be observed and any deviation from the continuum process must be carefully modeled until a final understanding comes out, with the advantage that the final state is not strongly interacting. Note that in this modeling the η meson will contribute to $\gamma\gamma \to \gamma\gamma$ in a small region of momentum [10], even so it has to be subtracted in order to extract the complete σ signal.



FIG. 2. Significance as a function of decay width into two photons, $\Gamma_{\gamma\gamma}$, for a sigma meson with mass equal to 478 MeV and total decay width of 324 MeV. The solid curve was obtained integrating the cross sections in the interval $438 < M < 519$ MeV, the dashed one in the interval $300 < M < 800$ MeV. The signal and background are giving by the dashed (Breit-Wigner) and solid (box diagram) curves of Fig.(1) respectively.

We changed the values of the σ mass and total width around the central ones reported by the E791 Collaboration. We do not observed large variations in our result, but noticed that it is quite sensitive to variations of the partial decay width into photons. It is interesting to look at the values of the significance which is written as $\mathcal{L}\sigma_{signal}/\sqrt{\mathcal{L}\sigma_{back}}$ and characterizes the statistical deviation of the number of the observed events from the predicted background. The significance as a function of the two photons decay width of the sigma meson, with mass equal to 478 MeV and total decay width of 324 MeV, is shown in Fig.(2), were we used a luminosity of $\mathcal{L} = 2.0 \times 10^{26}$ cm^{-2}s^{-1} at RHIC and assumed one year of operation. The significance is above 2σ 95% confidence level limit for two photon decay width greater than 4.7 keV, while for a 5σ discovery criteria can be obtained with $\Gamma_{\gamma\gamma} > 7.5$ keV. The numbers in Fig.(2) were computed with the signal given by the Breit-Wigner profile, and the background by the pure box diagram. The solid curve was obtained integrating the cross sections in the range of experimental mass uncertainty $438 < M < 519$ MeV, while the dashed curve resulted from the integration in the interval $300 < M < 800$ MeV. Note that there is no reason, a priori, to restrict the measurement to a small bin of energy. This choice will depend heavily on the experimental conditions. Therefore, for values of $\Gamma_{\gamma\gamma}$ already quoted in the literature the sigma meson has a chance to be seen in its two photon decay mode. The discovery limits discussed above refer only to a statistical evaluation. Our work shows the importance of the complete simulation of the signal and background including an analysis of possible systematic errors that may decrease the significance.

In summary, we discussed the possibility of observing the subprocess $\gamma\gamma \rightarrow \gamma\gamma$ in peripheral heavy ion collisions at RHIC. This process is important per se, but it is more interesting because it can be intermediated by a scalar (σ) resonance. The main advantage of this process is that the initial and final states are not strongly interacting, providing a much cleaner signal than the usually studied two pion decay of the σ meson. We computed the resonant process, the continuum one and discussed other possible background. The resonant was determined within the Breit-Wigner approximation and with an effective Lagrangian describing the σ interaction with photons. We determined a cut that leaves the cross sections apart by one order of magnitude when using a partial decay width of the sigma ($\Gamma_{\gamma\gamma}$) into two photons of 4.7 keV. Larger and smaller values for $\Gamma_{\gamma\gamma}$ can be found in the literature. The significance for the σ discovery in this reaction was determined and the observation is possible with one year of data acquisition. It is important to perform a complete simulation of the signal and background including an analysis of possible systematic errors that may occur at the RHIC detectors.

ACKNOWLEDGMENTS

We would like to thank A. C. Aguilar, I. Bediaga, C. Dib, M. M. Leite, R. Rosenfeld and A. Zimerman for many valuable discussions. This research was supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (AAN), by Fundacão de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (CGR,AAN) and by Programa de Apoio a Núcleos de Excelência (PRONEX).

---

[1] N. A. Tornqvist and M. Roos, *Phys. Rev. Lett.* **76**, 1575 (1996); M. R. Pennington, Talk at the Workshop on Hadron Spectroscopy (WHS99), Frascati (March, 1999), hep-ph/9905241; N. A. Tornqvist, Summary talk of the conference on the sigma resonance (Sigma-meson 2000), Kyoto (June, 2000), hep-ph/0008136.

[2] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122**, 345 (1961); **124**, 246 (1961).

[3] R. Machleidt, K. Holinde and Ch. Elster, *Phys. Rep.* **149**, 1 (1987); T. Hatsuda, T. Kunihiro and H. Shimizu, *Phys. Rev. Lett.* **82**, 2840 (1999); T. Kunihiro, Talk at the Workshop on Hadron Spectroscopy (WHS99), Frascati (March, 1999), hep-ph/9905262.

[4] D. E. Groom *et al.*, *Eur. Phys. J.* **C15**, 1 (2000).

[5] E791 Collaboration, E. M. Aitala et al., *Phys. Rev. Lett.* **86**, 770 (2001).

[6] F. E. Close, G. R. Farrar and Z. Li, *Phys. Rev.* **D55**, 5749 (1997).

[7] G. Baur, K. Hencken, D. Trautmann, S. Sadovsky and Y. Kharlov, hep-ph/0112211 *Phys. Rep., in press*; C. A. Bertulani and G. Baur, *Phys. Rep.* **163**, 299 (1988); G. Baur, J. Phys. **G24**, 1657 (1998); S. Klein and E. Scannapieco, hep-ph/9706358 (LBNL-40457); J. Nystrand and S. Klein, hep-ex/9711021 (LBNL-42524); C. A. Bertulani, nucl-th/0011065, nucl-th/0104059.

[8] G. Baur and C. A. Bertulani, *Nucl. Phys.* **A505**, 835 (1989).

[9] R. N. Cahn and J. D. Jackson, Phys. Rev. **D42**, 3690 (1990).

[10] A. A. Natale, C. G. Roldão and J. P. V. Carneiro, *Phys.Rev.* **C65**, 014902 (2002).

[11] R. Karplus and M. Neuman, *Phys. Rev.* **83**, 776 (1951).

[12] B. De Tollis, *Nuovo Cimento* **32**, 757 (1964); **35**, 1182 (1965).

[13] C. G. Roldão and A. A. Natale, *Phys. Rev.* **C61**, 064907 (2000).

[14] J. Nystrand and S. Klein (LBNL-41111), Talk at the Workshop on Photon Interactions and the Photon Structure, Lund (Sep., 1998), nucl-ex/9811007.

[15] S. Klein and J. Nystrand, *Phys. Rev.* **C60**, 014903 (1999)

[16] D. Alde et al., *Z. Phys.* **C36**, 603 (1987).

[17] M. Boglione and M. R. Pennington, *Eur. Phys. J.* **C9**, 11 (1999).

[18] F. Sannino and J. Schechter, *Phys. Rev.* **D52**, 96 (1995); M. Harada, F. Sannino and J. Schechter, *Phys. Rev.* **D54**, 1991 (1996); *Phys. Rev. Lett.* **78**, 1603 (1997).

[19] J. D. Jackson, *Nuovo Cimento* **34**, 1644 (1964).

[20] B. De Tollis and G. Violini, *Nuovo Cimento* **41A**, 12 (1966).